

Degree condition for the existence of a k -factor containing a given Hamiltonian cycle

Yunshu Gao^a, Guojun Li^{a,*}, Xuechao Li^b

^a School of Mathematics and System Science, Shandong University, Jinan, 250100, PR China

^b Division of Academic Enhancement, University of Georgia, USA

ARTICLE INFO

Article history:

Received 10 October 2006

Received in revised form 2 May 2008

Accepted 12 May 2008

Available online 24 June 2008

Keywords:

Factor

Degree condition

Hamiltonian cycle

ABSTRACT

Let k be an integer with $k \geq 2$ and let G be a graph having sufficiently large order n . Suppose that kn is even, the minimum degree of G is at least k and $\max\{d_G(x), d_G(y)\} \geq (n + \alpha)/2$ for each pair of nonadjacent vertices x and y in G , where $\alpha = 3$ for odd k and $\alpha = 4$ for even k . Then G has a k -factor (i.e. a k -regular spanning subgraph) which contains a given Hamiltonian cycle C if $G - E(C)$ is connected.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction and results

Graphs considered here are finite undirected graphs without loops and multiple edges. For notation and terminology not defined here we refer the reader to [1].

Let $G = (V, E)$ be a simple graph. We use $V(G)$, $E(G)$, $\delta(G)$ to denote the vertex set, edge set, *minimum degree* in G , respectively. The order of G is $|G| = |V| = n$ and its size is $e(G) = |E|$. A graph is said to be *Hamiltonian* if it contains a Hamiltonian cycle. For $x \in V(G)$, we denote by $N(x)$ the neighborhood of x in G , by $d(x)$ the degree of x in G . For $S \subseteq V(G)$, let $|S|$ denote the number of the vertices in S and $G[S]$ be the subgraph induced by S . In particular, we write $G - S$ for $G[V(G) \setminus S]$. Let H be a subgraph of G and $u \in V(G)$ a vertex, $N(u, H)$ is the set of neighbors of u contained in H . We write $d(u, H) = |N(u, H)|$. Clearly, $d(u, G)$ is the degree of u in G , and we write $d(x)$ to replace $d(x, G)$. If there is no fear of confusion, we often identify a subgraph H of G with its vertex set $V(H)$. For any two vertex-disjoint subgraphs of G , say G_1 and G_2 , the *join* G^* of these two subgraphs is defined as follows: $V(G^*) = V(G_1) \cup V(G_2)$ and $E(G^*) = E(G_1) \cup E(G_2) \cup \{xy | x \in V(G_1), y \in V(G_2)\}$. We use $G_1 \otimes G_2$ to denote the join of G_1 and G_2 . We define the distance $d(x, y)$ between two vertices x and y as the minimum of the lengths of the $x - y$ paths of G . Given a disjoint subset $A, B \subseteq V(G)$, we write $e_G(A, B)$ for the number of edges in G joining a vertex in A to that in B . For a positive integer k , a k -factor is a spanning subgraph F such that $d_F(x) = k$ for each $x \in V(G)$.

A classic sufficient degree condition for a Hamiltonian graph was obtained by Ore.

Theorem 1 (Ore [2]). *If $d(u) + d(v) \geq n$ for each pair of nonadjacent vertices $u, v \in V(G)$, then G is Hamiltonian.*

Fan (see [3]) extended Theorem 1 by providing an improved sufficient condition for a Hamiltonian graph.

* Corresponding author.

E-mail addresses: gysh2004@mail.sdu.edu.cn (Y. Gao), gjli@sdu.edu.cn (G. Li).

Theorem 2 (Fan [3]). Let G be a 2-connected graph of order $n \geq 3$. If for any two vertices x and y of G such that $d(x, y) = 2$, $\max\{d(x), d(y)\} \geq n/2$, then G has a Hamiltonian cycle.

Theorem 3 (Iida and Nishimura [4]). Let $k \geq 1$ be an integer and let G be a graph of order $n \geq 4k - 5$. Suppose that kn is even, $\delta(G) \geq k$, and $d(u) + d(v) \geq n$ for each pair of nonadjacent vertices $u, v \in V(G)$. Then G has a k -factor.

Theorem 4 (Nishimura [5]). Let G be a connected graph of order $n \geq 4k - 3$, where $k \geq 3$. Suppose that kn is even, $\delta(G) \geq k$, and $\max\{d(x), d(y)\} \geq n/2$ for each pair of nonadjacent vertices $u, v \in V(G)$. Then G has a k -factor.

Recently, H. Matsuda [6] proved the following result.

Theorem 5 (Matsuda [6]). Let $k \geq 2$ be an integer and let G be a graph of order $n > 8k^2 - 2(\alpha + 12)k + 3\alpha + 16$, where $\alpha = 3$ for odd k and $\alpha = 4$ for even k . Suppose that kn is even and the minimum degree $\delta(G)$ of G is at least k . If for any nonadjacent vertices x and y of G , $d(x) + d(y) \geq n + \alpha$, then G has a k -factor containing a given Hamiltonian cycle.

In this paper, we improve the result of Theorem 5 by the following Theorem:

Theorem 6. Let $k \geq 2$ be an integer and let G be a graph of order $n > 12(k - 2)^2 + 2(5 - \alpha)(k - 2) - \alpha$. Suppose that kn is even, $\delta(G) \geq k$ and $\max\{d(x), d(y)\} \geq (n + \alpha)/2$ for each pair of nonadjacent vertices x and y in G , where $\alpha = 3$ for odd k and $\alpha = 4$ for even k . Then G has a k -factor which contains a given Hamiltonian cycle C if $G - E(C)$ is connected.

2. Main result

We list Tutte's Theorem regarding the existence of a k -factor which will be used in our proof of the main result.

Theorem 7 (Tutte [7]). Let G be a graph and $k \geq 1$ an integer. Then G has a k -factor if and only if

$$\theta_G(S, T, k) = k|S| + \sum_{x \in T} (d_{G-S}(x) - k) - h_G(S, T, k) \geq 0$$

for all disjoint subsets S and T of $V(G)$, where $h_G(S, T, k)$ is the number of the components D of $G - (S \cup T)$ such that $k|D| + e_G(V(D), T) \equiv 1 \pmod{2}$. Furthermore, whether G has a k -factor or not, we have that $\theta_G(S, T, k) \equiv k|V(G)| \pmod{2}$ for any disjoint subsets S and T of $V(G)$.

We call such a component D *odd component*. Let $\omega(G)$ denote the number of components of G and let $o(G)$ denote the number of components of G each of which has odd order. We provide several propositions.

Proposition 1. If the condition of Theorem 6 is satisfied, $k \geq 3$ and $G - E(C)$ is connected for any Hamiltonian cycle C in G . Then $\omega(G - E(C) - A) \leq |A| + 1$ for every $A \subset V(G)$ or $k = 3$ and G has a 3-factor which contains C .

Proof. Let $H := G - E(C)$. Suppose there exists a set $A \subset V(G)$ such that $\omega(H - A) \geq |A| + 2$. If $A = \emptyset$, it is obvious. Hence, we may assume that $A \neq \emptyset$. Let $C_1, C_2, \dots, C_\omega$ be the components of $H - A$. We assume that $|C_1| \leq |C_2| \leq \dots \leq |C_\omega|$. Since H is connected and $\omega \geq 2$ by the assumption, there exists a vertex $x_i \in V(C_i)$ with $N(x_i) \cap A \neq \emptyset$ for every $i \in \{1, 2, \dots, \omega\}$.

We claim that there exist $x_j, x_l \in \{x_1, x_2, \dots, x_{|A|+1}\}$ such that x_j and x_l are nonadjacent. Otherwise, we may assume that all the vertices of $V(C_1)$ are adjacent to those of $V(C_2)$ in C and $|C_1| \leq |C_2| \leq 2$. Suppose that $|A| \geq 2$, then $\omega(H - A) \geq 4$. If $|C_2| = 2$, then for each $u_1 \in V(C_1)$, $e_C(u_1, V(C_2)) = 2$ and so $u_1 u_3 \notin E(G)$ for each $u_3 \in V(C_3)$. If $|C_2| = 1$, then $u_1 u_2 \in E(G)$ for $\{u_1\} = V(C_1)$ and $\{u_2\} = V(C_2)$. Consequently, there exists $u_3 \in V(C_3)$ such that $u_1 u_3 \notin E(C)$ or $u_2 u_3 \notin E(C)$ and we are done. Hence, it remains the case that $|A| = 1$ and $\omega(H - A) = 3$. Clearly, $|C_1| = 1$ and $|C_2| \leq 2$, denote $A = \{z\}$. If $|C_2| = 1$, then $V(C_2) = \{x_2\}$. By the symmetry role of x_1 and x_2 , there exist two types of configurations of G along C , see Figs. 1 and 2, where $V(C_3) = V(P) \cup \{u\}$. If $|C_2| = 2$, denote $V(C_2) = \{x_2, x'_2\}$. Consequently, there exist three types of configurations according to whether $zx'_2 \in E(G)$ or $zx'_2 \notin E(G)$, where $V(C_3) = V(P) \cup \{u\}$, see Fig. 3. Note that in either case, $k = 3$ since $d_G(x_1) = 3$, then $\alpha = 3$. Furthermore, Type 2 will not appear. As n is even, then $\max\{d_H(w_1), d_H(w_2)\} \geq \frac{n}{2}$ for each pair of nonadjacent vertices $w_1, w_2 \in V(H)$, applying Theorem 2, $G - E(C)$ contains a Hamiltonian cycle and so G contains two edge-disjoint Hamiltonian cycles. Consequently, $G - E(C)$ contains a 1-factor since n is even, denoted by \mathcal{F} . Then, $E(C) \cup \mathcal{F}$ forms a desired 3-factor which contains C .

From the above claim, by the condition of Theorem 6, at least one of the two vertices, say x_j , has degree at least $\frac{n+\alpha-4}{2}$ in H . This yields

$$\frac{n + \alpha - 4}{2} \leq d_H(x_j) \leq |V(C_j)| - 1 + |A|.$$

Therefore, for every $i \geq j$,

$$|V(C_i)| \geq \frac{n + \alpha - 4}{2} - |A| + 1.$$




Fig. 1. Type 1.





Fig. 2. Type 2.

Fig. 3. Three configurations along C .

Since $j \leq |A| + 1$ and $3 \leq \alpha \leq 4$, we obtain

$$n \geq |A| + \sum_{i=1}^{|A|} V(C_i) + \sum_{i=|A|+1}^{|A|+2} V(C_i) \geq 2|A| + 2 \left(\frac{n + \alpha - 4}{2} - |A| + 1 \right) = n + 1,$$

a contradiction. \square

Proposition 2. Let G be a graph satisfying the condition of Theorem 6. Suppose that for each Hamiltonian cycle C in G , $H = G - E(C)$ is connected and $k \geq 3$. Then G has a 3-factor which contains any given Hamiltonian cycle C .

Proof. Suppose for a contradiction, for a given Hamiltonian C in G , G does not contain a 3-factor which contains C , this implies $G - E(C)$ does not contain a 1-factor. By Tutte's 1-factor Theorem [8], there exists a set $A \subset V(G)$ such that $o(H - A) > |A|$. Since $3n$ is even by the condition of Theorem 6, then n is even. It follows that $o(H - A) \geq |A| + 2$, which contradicts Proposition 1. \square

Proposition 3. Let G be a graph satisfying the condition of Theorem 6. Suppose that for each Hamiltonian cycle C in G , $H = G - E(C)$ is connected and $k \geq 3$. Then $\omega(H - A) \leq |A| + 5 - \alpha$ for all $\emptyset \neq A \subseteq V(H)$ or $k = 3$ and G has a 3-factor which contains C .

Proof. It is obvious by applying Proposition 1. \square

Proof of Theorem 6. We prove it by contradiction. We may assume that $k \geq 3$ since G contains a Hamiltonian cycle by Theorem 2. Let $H := G - E(C)$, $\rho := k - 2$ where $k \geq 3$. In the following, we may assume that for each Hamiltonian C ,

$G - E(C)$ is connected. Clearly, $V(H) = V(G)$ and $\rho \geq 1$,

$$d_H(x) = d(x) - 2 \geq \rho \quad \text{for all } x \in V(H), \quad (1)$$

and since C is a Hamiltonian cycle of G , we have

$$\max\{d_H(u), d_H(v)\} \geq \frac{n + \alpha - 4}{2} \quad \text{for each nonadjacent } u, v \in V(H). \quad (2)$$

Obviously, G has the desired factor if and only if H has a ρ -factor. Suppose to the contrary, H does not contain such a factor, then by [Theorem 7](#), there exist disjoint subsets S and T of $V(H)$ such that

$$\theta_H(S, T, \rho) = \rho|S| + \sum_{x \in T} (d_{H-S}(x) - \rho) - h_H(S, T, \rho) \leq -2. \quad (3)$$

We choose such subsets S and T such that $|T|$ is minimum whereas $|S|$ is maximal with respect to $|T|$. Further, we choose ρ as small as possible.

In view of [Proposition 2](#), we see that H contains 1-factor. So in the following proof we assume that $\rho \geq 2$.

Claim 1. $|T| \geq |S| + 1$.

Proof. We first show that H has a $(\rho - 2)$ -factor. It is trivial if $\rho = 2$. So we assume that $\rho \geq 3$. By the choice of ρ , H has a $(\rho - 2)$ -factor, by [Theorem 7](#), we have $\theta_H(S, T, \rho - 2) \geq 0$. Note that $h_H(S, T, \rho) = h_H(S, T, \rho - 2)$ by the definition of odd components, we obtain $-2 \geq \theta_H(S, T, \rho) - \theta_H(S, T, \rho - 2) = 2|S| - 2|T|$, which implies that $|T| \geq |S| + 1$. \square

By [Claim 1](#), it is clear that $T \neq \emptyset$.

Claim 2. $2|S| \leq n - 6\rho + \alpha - 3$.

Proof. Suppose that $2|S| \geq n - 6\rho + \alpha - 2$, that is, $n - 2|S| \leq 6\rho - \alpha + 2$. Thus $|T| - |S| = n - 2|S| - |V(H) - (S \cup T)| \leq 6\rho - \alpha + 2 - h_H(S, T, \rho)$. By (3), we obtain that

$$\begin{aligned} \sum_{x \in T} d_{H-S}(x) &\leq \rho|T| - \rho|S| + h_H(S, T, \rho) - 2 \\ &\leq \rho(|T| - |S|) + h_H(S, T, \rho) - 2 \\ &\leq \rho(6\rho - \alpha + 2 - h_H(S, T, \rho)) + h_H(S, T, \rho) - 2 \\ &\leq \rho(6\rho - \alpha + 2) - 2. \end{aligned}$$

Since $n > 12\rho^2 + 2(5 - \alpha)\rho - \alpha$, we have $|T| \geq |S| + 1 \geq \frac{n}{2} - 2\rho + \frac{\alpha}{2} > 6\rho^2 + (3 - \alpha)\rho > 2$. Consequently, together with [Claim 1](#), we have

$$\begin{aligned} \frac{\sum_{x \in T} d_{H-S}(x)}{|T| - 2} &\leq \frac{\rho(6\rho - \alpha + 2) - 2}{|T| - 2} \\ &\leq \frac{\rho(6\rho - \alpha + 2) - 2}{|S| - 1} \\ &\leq \frac{2\rho(6\rho - \alpha + 2) - 4}{n - 6\rho + \alpha - 4} < 1, \end{aligned}$$

where the last estimation follows from $n > 12\rho^2 + 2(5 - \alpha)\rho - \alpha$. It follows that

$$\sum_{x \in T} d_{H-S}(x) \leq |T| - 3. \quad (4)$$

Let $T_0 = \{x \in T \mid d_{H-S}(x) = 0\}$. It follows from [Claim 1](#) that $n \geq |S| + |T| \geq 2|S| + 1$. Hence if n is even, $n \geq 2|S| + 2$. By the definition of α and the assumption that kn is even, $n \geq 2|S| + 2$ if $\alpha = 3$ and $n \geq 2|S| + 1$ if $\alpha = 4$. In either case, we have $d_H(x) \leq |S| < \frac{n + \alpha - 4}{2}$ for any $x \in T_0$. Note that $|T_0| \geq 3$ by inequality (4), there exist two vertices $x, y \in T_0$ such that $xy \notin E(G)$. Hence, we obtain that

$$\frac{n + \alpha}{2} \leq \max\{d(x), d(y)\} \leq |S| + |N_C(x)| < \frac{n + \alpha - 4}{2} + 2 < \frac{n + \alpha}{2},$$

which contradicts [Claim 1](#). \square

Claim 3 (*H. Matsuda, [6]*). For any $x \in T$, $d_{H-S}(x) \leq \rho - 2 + h_H(S, T, \rho)$ and $d_T(x) \leq \rho - 2$.

Proof. Let $T' = T \setminus \{x\}$ for any $x \in T$. By the choice of T we have $\theta_H(S, T', \rho) \geq 0$. Since $\theta_H(S, T, \rho) \leq -2$, we obtain $2 \leq \theta_H(S, T', \rho) - \theta_H(S, T, \rho) \leq \rho - d_{H-S}(x) + h_H(S, T, \rho) - h_H(S, T', \rho)$ which implies $d_{H-S}(x) \leq \rho - 2 + h_H(S, T, \rho) - h_H(S, T', \rho)$. This inequality, together with $e_H(x, H - (S \cup T)) \geq h_H(S, T, \rho) - h_H(S, T', \rho)$ yields $d_T(x) = d_{H-S}(x) - e_H(x, H - (S \cup T)) \leq \rho - 2$. \square

Claim 4 (H. Matsuda, [6]). $e_H(u, T) \leq \rho - 2$ for any $u \in V(H) \setminus (S \cup T)$.

Proof. By the maximality of S , $\theta_H(S \cup \{u\}, T, \rho) \geq 0$ for any $u \in V(H) \setminus (S \cup T)$. Note that $\theta_H(S, T, \rho) \leq -2$, we have $2 \leq \theta_H(S \cup \{u\}, T, \rho) - \theta_H(S, T, \rho) \leq \rho - e_H(u, T)$, which implies that $e_H(u, T) \leq \rho - 2$. \square

For the sake of convenience, let $C_1, C_2, \dots, C_\omega$ be the odd components of $H - (S \cup T)$ with $|C_1| \leq \dots \leq |C_\omega|$, where $\omega = h_H(S, T, \rho)$ is the number of odd components of $H - (S \cup T)$.

Claim 5. For $x \in V(C_i)$, if $d(x) \geq \frac{n+\alpha}{2}$, then $2|C_i| \geq n - 2|S| - 2\rho + \alpha + 2$.

Proof. By Claim 4, we have

$$\begin{aligned} \frac{n+\alpha}{2} \leq d(x) &\leq e_H(x, T) + |S| + |C_i| - 1 + |N_C(x)| \\ &\leq \rho + |S| + |C_i| - 1. \end{aligned}$$

Hence $2|C_i| \geq n - 2|S| - 2\rho + \alpha + 2$. \square

Claim 6. $\omega = h_H(S, T, \rho) \leq 4$.

Proof. Suppose that $\omega = h_H(S, T, \rho) \geq 5$. There are no two nonadjacent vertices $u_1 \in V(C_1)$ and $u_2 \in V(C_2)$. Otherwise by the assumption of G , we assume that $d(u_i) \geq \frac{n+\alpha}{2}$ where $i = 1$ or 2 . So $2|C_i| \geq n - 2|S| - 2\rho + \alpha + 2$ ($i = 1$ or 2) by Claim 5. By Claim 2, we obtain

$$\begin{aligned} n &\geq |S| + |T| + |C_1| + |C_2| + |C_3| + |C_4| + |C_5| \\ &\geq 2|S| + 1 + (6-i)|C_i| \\ &\geq 2|S| + 1 + 2(n - 2|S| - 2\rho + \alpha + 2) \\ &> n + 1, \end{aligned}$$

a contradiction. Thus each vertex of $V(C_1)$ is adjacent to each vertex of $V(C_2)$ in G and so is in C . Then we have $|C_1| \leq |C_2| \leq 2$. If $|C_2| = 2$, then for each $u_1 \in V(C_1)$, $e_C(u_1, V(C_2)) = 2$ and thus $u_1 u_3 \notin E(C)$ for each vertex $u_3 \in V(C_3)$. If $|C_2| = 1$, then $u_1 u_2 \in E(C)$ for $\{u_1\} = V(C_1)$ and $\{u_2\} = V(C_2)$. Thus there exists a vertex $u_3 \in V(C_3)$ such that at least one of $\{u_1 u_3, u_2 u_3\}$ is not in $E(C)$. We assume that $u_1 u_3 \notin E(C)$. If $d(u_1) \geq \frac{n+\alpha}{2}$, then it is the previous case. If $d(u_3) \geq \frac{n+\alpha}{2}$, by Claim 5, $2|C_3| \geq n - 2|S| - 2\rho + \alpha + 2$. This inequality together with Claim 2 gives

$$\begin{aligned} n &\geq |S| + |T| + |C_1| + |C_2| + |C_3| + |C_4| + |C_5| \\ &\geq |S| + |T| + 2 + 3|C_3| \\ &\geq 2|S| + 1 + 2 + \frac{3}{2} \times (n - 2|S| - 2\rho + \alpha + 2) \\ &> n + 1, \end{aligned}$$

a contradiction again. \square

Claim 7. $G[T]$ is a complete subgraph of G .

Proof. For $xy \notin E(G)$ where $x, y \in V(T)$, we may assume $d(x) \geq \frac{n+\alpha}{2}$, then by Claims 3 and 6,

$$\frac{n+\alpha}{2} \leq d(x) \leq e(x, S) + d_{H-S}(x) + 2 \leq |S| + \rho - 2 + h_H(S, T, \rho) + 2 \leq \rho + 4 + |S|.$$

Since $\rho \geq 2$, according to Claim 2, $n + \alpha \leq 2(\rho + 4 + |S|) \leq 2\rho + 8 + n - 6\rho + \alpha - 3 = n + \alpha - 4\rho + 5 < n + \alpha$, a contradiction. Thus $xy \in E(G)$. \square

Define $m_1 = \min\{d_{H-S}(x) | x \in T\}$ and let $x_1 \in T$ be a vertex with $d_{H-S}(x_1) = m_1$. If $m_1 \geq \rho + 2$, since $|T| \geq 1$, then by Proposition 3, $\theta_H(S, T, \rho) \geq \rho|S| + \sum_{x \in T} (m_1 - \rho) - h_H(S, T, \rho) \geq |S| + |T| - \omega(H - (S \cup T)) + |T| \geq |T| - 2 \geq -1$, which contradicts inequality (3), hence $m_1 \leq \rho + 1$.

By Claims 1, 3 and 7 we have $|S| + 1 \leq |T| \leq d_T(x_1) + |\{x_1\}| + |N_C(x_1)| \leq \rho + 1$ which implies that $|S| \leq \rho$ and $|T| \leq \rho + 1$.

Claim 8. $|C_1| \geq 2$.

Proof. Suppose that $|C_1| = |\{u\}| = 1$. By Claim 4, $e_H(u, T) \leq \rho - 2$. If $e_H(u, T) = \rho - 2$, then $\rho + e_H(u, T) = 2\rho - 2$. This contradicts the fact that C_1 is an odd component of $H - (S \cup T)$, thus we assume $e_H(u, T) \leq \rho - 3$. By (1), $|S| + e_H(u, T) \geq d_H(u) \geq \delta(H) \geq \rho$, then $|S| \geq 3$. If $|T| \leq \rho - 1$, by Claim 6, inequality (3) and the fact $|S| + d_{H-S}(x) \geq \delta(H) \geq \rho$ for every $x \in T$, we have $-2 \geq \theta_H(S, T, \rho) \geq |S| + \sum_{x \in T} (|S| + d_{H-S}(x) - \rho) - h_H(S, T, \rho) \geq |S| - h_H(S, T, \rho) \geq -1$. Hence, a contradiction. So we assume that $|T| \geq \rho$. So $|T| = \rho$ or $\rho + 1$. By Claim 7, we obtain $d_{H-S}(x) \geq d_T(x) \geq \rho - 3$ for each $x \in T$, implying $\rho \geq 3$. If $e_H(u, T) \leq \rho - 4$, then $|S| \geq 4$. Since $G[T]$ is complete by Claim 7, $|E(G[T])| = |T|(|T| - 1)/2$. As C is a Hamiltonian cycle, $|E(G[T]) \cap C| \leq |T| - 1$. Consequently, we obtain that

$$\begin{aligned} \sum_{x \in T} d_{H-S}(x) &\geq 2|E(G[T]) \setminus E(C)| \\ &\geq |T|(|T| - 1) - 2(|T| - 1) = (|T| - 1)(|T| - 2). \end{aligned}$$

Then it follows from Claim 6 and the fact $|S| \geq 4$ that

$$\begin{aligned} \theta_H(S, T, \rho) &= \rho|S| + \sum_{x \in T} (d_{H-S}(x) - \rho) - h_H(S, T, \rho) \\ &\geq \rho|S| + (|T| - 1)(|T| - 2) - \rho|T| - h_H(S, T, \rho) \\ &\geq 4\rho + |T|^2 - (\rho + 3)|T| - 2 \\ &= (|T| - 3)(|T| - \rho) + \rho - 2. \end{aligned} \quad (5)$$

Note that $\rho \geq 3$, if $|T| = \rho$, then $\theta_H(S, T, \rho) \geq 1$, which contradicts (3). Hence, $|T| = \rho + 1$, then $|T| \geq 3$ and by (5), we see that $\theta_H(S, T, \rho) \geq 1$, again a contradiction. Thus, we have $e_H(u, T) = \rho - 3$. If $|T| = \rho + 1$, then by Claim 7, $d_{H-S}(x) \geq \rho - 2$ for each $x \in T$. Since $|E(G[T]) \cap C| \leq |T| - 1$, then $\theta_H(S, T, \rho) \geq 3\rho - (\rho - 1) - 8 = 2\rho - 7 \geq -1$, which contradicts (3). Hence, it remains the case that $|T| = \rho$, by Claim 7, $d_{H-S}(x) \geq \rho - 3$ for each $x \in T$. Recalling that H is connected and $\rho \geq 3$, therefore, we have $\theta_H(S, T, \rho) \geq 3\rho + (\rho - 3)(\rho - 2 - \rho) - 10 = \rho - 4 \geq -1$, which contradicts (3). Hence we complete the proof of Claim 8. \square

By Claim 6, $h_H(S, T, \rho) \leq 4$. We will finish the proof by using the following four Claims.

Claim A. $h_H(S, T, \rho) \neq 4$.

Proof. Otherwise, by Claim 8, $2 \leq |C_1| \leq |C_2| \leq |C_3| \leq |C_4|$, thus there exist two vertices $u_1 \in V(C_1)$ and $u_2 \in V(C_2)$ such that $u_1 u_2 \notin E(G)$. If $d(u_1) \geq \frac{n+\alpha}{2}$, then by Claims 1, 2 and 5, we have

$$\begin{aligned} n &\geq |S| + |T| + |C_1| + |C_2| + |C_3| + |C_4| \geq 2|S| + 1 + 4|C_1| \\ &\geq 2|S| + 1 + 2(n - 2|S| - 2\rho + \alpha + 2) \\ &> n + 1. \end{aligned}$$

This is a contradiction. Therefore, $d(u_1) < \frac{n+\alpha}{2}$, which implies $d(u_2) \geq \frac{n+\alpha}{2}$, then $2|C_2| \geq n - 2|S| - 2\rho + \alpha + 2$ by Claim 5. This inequality, together with Claims 1, 2 and 8, gives that

$$\begin{aligned} n &\geq |S| + |T| + |C_1| + |C_2| + |C_3| + |C_4| \\ &\geq 2|S| + 1 + 2 + \frac{3}{2} \times (n - 2|S| - 2\rho + \alpha + 2) \\ &> n + 1, \end{aligned}$$

a contradiction. \square

Claim B. $h_H(S, T, \rho) \neq 2$.

Proof. We prove it by contradiction. Assume that $h_H(S, T, \rho) = 2$. First we show that $m_1 = \min\{d_{H-S}(x) | x \in T\} \leq \rho$. Otherwise, $m_1 = \rho + 1$, and by (3), $2 = h_H(S, T, \rho) \geq \rho|S| + (m_1 - \rho)|T| + 2$ which implies that $S \cup T = \emptyset$, a contradiction. Therefore, $m_1 \leq \rho$. Since $|S| + d_{H-S}(x_1) \geq \delta(H) \geq \rho$, then $|S| \geq \rho - m_1$. This inequality, together with (3), gives us that

$$\begin{aligned} 2 = h_H(S, T, \rho) &\geq \rho|S| + (m_1 - \rho)|T| + 2 \\ &\geq \rho(\rho - m_1) + (m_1 - \rho)|T| + 2 \\ &= (\rho - m_1)(\rho - |T|) + 2 \geq 2. \end{aligned} \quad (6)$$

Since $\rho \geq m_1$, the last inequality holds when $|T| \leq \rho$ unless $|T| = \rho + 1$. When $|T| = \rho + 1$, by Claim 7, $m_1 \geq |T| - 3 = \rho - 2$. Clearly, $m_1 \leq \rho - 1$. If $|S| = 0$, then $m_1 = \rho$, then inequality (6) still holds. So, $|S| \geq 1$. If $|S| = 1$, then $m_1 \geq \rho - 1$. By Claim 8 and the minimality of T , we have $\theta_H(S, T, \rho) \geq \rho + (\rho + 1 - 2)(\rho - 1 - \rho) + 2(\rho - \rho) - 2 = \rho - \rho - 1 = -1$, which contradicts (3). Therefore, $|S| \geq 2$. By Claim 8 and the minimality of T , it follows that $\theta_H(S, T, \rho) \geq 2\rho + (\rho + 1 - 3)(\rho - 2 - \rho) + 3(\rho - 1 - \rho) - 2 = -1$, which contradicts (3) again.

Consequently, it follows from (6) that $(\rho - m_1)(\rho - |T|) = 0$. This implies that $|S| + m_1 = \rho$ and $\rho = m_1$ or $\rho = |T|$. Note that $|T| \leq m_1 + 1 \leq \rho + 1$, $|T| = \rho$ implies that $m_1 = \rho - 1$ or ρ , then $|S| = 1$ or $|S| = 0$. Next we consider the following two cases.

Case (B-1). $|S| = 1$, $|T| = \rho$ and $d_{H-S}(x) = \rho - 1$ for any $x \in T$.

In this case, $\frac{n+\alpha}{2} > d(x) \geq \rho - 1 + |S| + |N_C(x)|$ for any $x \in T$. Since $G[T]$ is a complete subgraph by Claim 7, we have $e_G(T, H - (S \cup T)) = 0$, therefore, for each $u_1 \in V(C_1)$, we obtain

$$\begin{aligned} d(u_1) &\leq \frac{n - |S| - |T|}{2} - 1 + |S| + |N_C(u_1)| \\ &= \frac{n + 3 - \rho}{2} < \frac{n + \alpha}{2}. \end{aligned}$$

Consequently, we find a pair of nonadjacent vertices $u_1 \in V(C_1)$ and $x \in G[T]$ such that $\max\{d(u_1), d(x)\} < \frac{n+\alpha}{2}$, a contradiction.

Case (B-2). $|S| = 0$, $m_1 = \rho$, $|T| = \rho$ and $d_{H-S}(x) = \rho$ for any $x \in T$.

Note $\frac{n+\alpha}{2} > d(x) \geq \rho - 1 + |S| + |N_C(x)|$ for any $x \in T$. We have $\theta_H(S, T, \rho) = \rho|S| + \sum_{x \in T} (d_{H-S}(x) - \rho) - 2 = -2$.

Since $m_1 = \rho$, then $\frac{n+\alpha}{2} > d(x_1) = m_1 + |N_C(x_1)| = \rho + 2$. On the other hand, by Claim 7, (3) and the fact $d_{H-S}(x) = \rho$ for each $x \in T$, there is at most one vertex u_i of C_i ($i = 1$ or 2) that satisfies $u_i \in N_H(x)$ ($x \in T$). Since $|C_1| \geq 2$ by Claim 8, we can find another vertex $y \in V(C_1)$ such that $xy \notin E(H)$. Similarly as Case (B-1), we obtain

$$\begin{aligned} d_H(y) &\leq \frac{n - |S| - |T|}{2} - 1 + |S| \\ &= \frac{n - 2 - \rho}{2} < \frac{n + \alpha - 4}{2} \end{aligned}$$

which contradicts the inequality (2). \square

Claim C. $h_H(S, T, \rho) \neq 3$.

Proof. Otherwise, by Claim 8, $2 \leq |C_1| \leq |C_2| \leq |C_3|$, thus there exist two vertices $u_1 \in V(C_1)$ and $u'_2 \in V(C_2)$ such that $u_1 u'_2 \notin E(G)$. If $d(u_1) \geq \frac{n+\alpha}{2}$, similarly, by Claims 1 and 2, we have

$$\begin{aligned} n &\geq |S| + |T| + |C_1| + |C_2| + |C_3| \geq |S| + |T| + 3|C_1| \\ &\geq 2|S| + 1 + \frac{3}{2} \times (n - 2|S| - 2\rho + \alpha + 2) \\ &> n + 1. \end{aligned} \quad (7)$$

This is a contradiction. So $d(u_1) < \frac{n+\alpha}{2}$, then $d(u'_2) \geq \frac{n+\alpha}{2}$. We have $2|C_2| \geq n - 2|S| - 2\rho + \alpha + 2$ by Claim 5. In fact, for any $x \in V(C_1)$, we have $d(x) < \frac{n+\alpha}{2}$ by using the same argument. If there exists $u_1 \in V(C_1)$ such that $u_1 x_1 \notin E(G)$, where $d_{H-S}(x_1) = m_1$ and $x_1 \in T$. By the degree condition, $\frac{n+\alpha}{2} \leq d(x_1) \leq |S| + m_1 + 2 \leq 2\rho + 3$ (since $m_1 \leq \rho + 1$), then $n \leq 2(2\rho + 3) - \alpha \leq 4\rho^2 + 3\rho - \alpha$, this inequality contradicts the fact that $n > 12\rho^2 + 2(5 - \alpha)\rho - \alpha$, thus $T \cup C_1 \subseteq N_G(x_1) \cup \{x_1\}$, which gives us that

$$|T \cup C_1| \leq |N_G(x_1) \cup \{x_1\}| \leq m_1 + 1 + 2 \leq \rho + 2 + 2 = \rho + 4. \quad (8)$$

Since $d_{G-S}(x) \leq d_{H-S}(x) + 2 \leq \rho + 3$ for all $x \in T$. Note $|S| \leq \rho$, $|T \cup C_1| \leq \rho + 4$, $\rho \geq 2$, $|T| \leq \rho + 1$ and $n > 12\rho^2 + 2(5 - \alpha)\rho - \alpha$, we have $n - |S \cup T \cup C_1| > (\rho + 3)(\rho + 1) + 1 \geq \sum_{x \in T} d_{G-S}(x) + 1$. The previous inequality implies that there exists a vertex $u_2 \in V(G) \setminus \{S \cup T \cup C_1\}$ such that $e_G(u_2, T) = 0$. Note that $\rho \geq 2$, by Claim 7 and (8), $d(x) = |T| - 1 \leq \rho + 3 < \frac{n+\alpha}{2}$ for each $x \in T$. Thus we may assume that $\frac{n+\alpha}{2} \leq d(u_2) \leq (|C_m| - 1 + |S| + 2)$ which leads us to that $2|C_m| \geq n + \alpha - 2 - 2|S|$, where C_m is the component such that $u_2 \in V(C_m)$. If $m \neq 2, 3$, then Claims 1 and 2 together with Claim 8 lead to

$$\begin{aligned} n &\geq |S| + |T| + |C_1| + |C_2| + |C_3| + |C_m| \\ &\geq 2|S| + 1 + 2 + (n - 2|S| - 2\rho + \alpha + 2) + \frac{n + \alpha - 2|S| - 2}{2} \\ &= 2|S| + 1 + 2 + n - 2|S| + \rho + \alpha + 2 + \frac{1}{2} \\ &> n + 1. \end{aligned}$$

This contradiction implies that $m = 2$ or 3 . In the following, we prove that $|S| = 0$. Otherwise, $|S| \geq 1$. If $m_1 = \rho$ or $\rho + 1$, then it follows that $\theta_H(S, T, \rho) \geq \rho|S| + (m_1 - \rho)|T| - h_H(S, T, \rho) \geq \rho - 3 \geq -1$ since $\rho \geq 2$. This contradicts (3). Thus

$m_1 \leq \rho - 1$, by $T \cup C_1 \subseteq N_G(x_1) \cup \{x_1\}$, we have

$$\begin{aligned} |T| + 2 &\leq |T \cup C_1| \leq d_{H-S}(x_1) + |\{x_1\}| + |N_C(x_1) \cap (T \cup C_1)| \\ &\leq m_1 + 1 + 2 \leq \rho + 2 \end{aligned} \quad (9)$$

which implies that $|T| \leq \rho$. We first assume that $|T| = \rho$. Then $|C_1| = 2$, $m_1 = \rho - 1$ and $|N_C(x_1) \cap (T \cup C_1)| = 2$. If $|S| \geq 2$, then $\theta_H(S, T, \rho) \geq \rho - 3 \geq -1$, a contradiction. So $|S| = 1$, then using a similar argument as in Case (B-1), we arrive at a contradiction to (2). Hence it remains the case that $|T| \leq \rho - 1$. If $|T| \leq \rho - 2$, then $-2 \geq \theta_H(S, T, \rho) = \rho|S| + \sum_{x \in T} (d_{H-S}(x) - \rho) - h_H(S, T, \rho) \geq 2|S| + \sum_{x \in T} (|S| + d_{H-S}(x) - \rho) - h_H(S, T, \rho) \geq 2|S| - h_H(S, T, \rho) \geq 2|S| - 3 \geq -1$ since $|S| + d_{H-S}(x) - \rho \geq \delta(H) - \rho \geq 0$, which contradicts (3). Hence $|T| = \rho - 1$. Then $-2 \geq \theta_H(S, T, \rho) = |S| + \sum_{x \in T} (|S| + d_{H-S}(x) - \rho) - h_H(S, T, \rho) \geq \sum_{x \in T} (|S| + d_{H-S}(x) - \rho) + |S| - 3 \geq |S| - 3 \geq -2$ which implies that for any $x \in T$, $|S| + d_{H-S}(x) = \rho$ and $|S| = 1$. Particularly, $m_1 = d_{H-S}(x_1) = \rho - |S| = \rho - 1$. By a similar argument as in Case (B-1), we obtain a contradiction.

Without loss of generality, suppose that $u_2 \in V(C_3)$. Since $S = \emptyset$, it follows that $e_G(x, T) > 0$ for any $x \in V(C_3)$. On the other hand, by Claim 3, for any $x \in T$, we have $d_H(x) \leq \rho + 1$. Note that $|C_3| \geq \frac{n-2\rho+\alpha+2}{2} > \rho + 1$, $|C_1| \geq 2$ and $T \cup C_1 \subseteq N_G(x_1) \cup \{x_1\}$, then there exists a vertex $v \in C_3$ such that $e_G(v, T) = 0$. This contradicts $e_G(x, T) > 0$ for any $x \in V(C_3)$. Hence we complete the proof of Claim C. \square

Claim D. $h_H(S, T, \rho) \not\leq 1$.

Proof. Otherwise, by (3), we have

$$-1 \geq h_H(S, T, \rho) - 2 \geq \rho|S| + \sum_{x \in T} (d_{H-S}(x) - \rho). \quad (10)$$

If $|T| \leq \rho$, then it follows from (10) that $-1 \geq \rho|S| + \sum_{x \in T} (d_{H-S}(x) - \rho) \geq \sum_{x \in T} (|S| + d_{H-S}(x) - \rho) \geq 0$, a contradiction. Thus we may assume that $|T| = \rho + 1$. If $m_1 \geq \rho$, then by (10), $-1 \geq \rho|S| + \sum_{x \in T} (d_{H-S}(x) - \rho) \geq \rho|S|$, again a contradiction. So $m_1 \leq \rho - 1$. Note that $|S| + d_{H-S}(x_1) \geq \delta(H) \geq \rho$, then $|S| \geq \rho - m_1 \geq 1$. Suppose that $|S| = 1$, then $m_1 = \rho - 1$, by the definition of m_1 , we have $d_{H-S}(x) \geq \rho - 1$ for each $x \in V(T)$. Note that $|E(G[T]) \cap C| \leq |T| - 1$, therefore, $\theta_H(S, T, \rho) \geq \rho + (\rho + 1 - 1)(\rho - 1 - \rho) + (\rho - \rho) - 1 = -1$, which contradicts (3). Thus we may assume $|S| \geq 2$. Since $G[T]$ is complete, we have $|E(G[T])| = |T|(|T| - 1)/2$. Note that C is a Hamiltonian cycle of G , $|E(G[T]) \cap C| \leq |T| - 1$. We obtain

$$\begin{aligned} \sum_{x \in T} d_{H-S}(x) &\geq 2|E(G[T])| - |E(C)| \\ &\geq |T|(|T| - 1) - 2(|T| - 1) = (|T| - 1)(|T| - 2). \end{aligned}$$

Noting that $|S| \geq 2$ and $|T| = \rho + 1$, we have

$$\theta_H(S, T, \rho) \geq \rho|S| + (|T| - 1)(|T| - 2) - \rho|T| - h_H(S, T, \rho) \geq 2\rho - 2\rho - 1 = -1$$

which contradicts (3).

Now, we complete the proof of Theorem 6. \square

3. Concluding remarks

Remark 1. Note that Theorem 4 cannot show the existence of a regular factor in $G - E(C)$. Consider a graph $G = K_t \oplus K_{2\alpha} \oplus K_{t+2\alpha}$ which is used in [6] where t is a sufficiently large integer. Then G satisfies the Ore-type condition and $G - E(C)$ is connected for any Hamiltonian cycle C . Note that $\delta_{G-E(C)} \geq k - 2$. For each $x \in V(G)$, we have

$$d(x) = \begin{cases} t + 2\alpha - 1, & \text{if } x \in V(K_t), \\ n - 1, & \text{if } x \in V(K_{2\alpha}), \\ t + 4\alpha - 1, & \text{if } x \in V(K_{t+2\alpha}). \end{cases}$$

However, $G - E(C)$ has two vertices $x, y \in V(K_t)$ such that

$$d_{G-E(C)}(x) + d_{G-E(C)}(y) = 2(|K_t| + |K_{2\alpha}|) - 2 - 4 = 2(t + 2\alpha) - 6 = n - 6.$$

Hence we cannot apply Theorem 3 to $G - E(C)$ to ensure the existence of $(k - 2)$ -factor. In particular, for any two nonadjacent vertices x and y of $V(K_t)$, $\max\{d_{G-E(C)}(x), d_{G-E(C)}(y)\} < \frac{n}{2}$. Consequently, we cannot apply Theorem 4 to $G - E(C)$ to ensure the existence of $(k - 2)$ -factor. Note that G satisfies the condition of Theorem 6. So our theorem (Theorem 6) guarantees that G has a k -factor but Theorems 3 and 4 do not.

Remark 2. In the proof of [Theorem 6](#), for each Hamiltonian cycle C in G , the requirement of connectedness of $G - E(C)$ is necessary. For example, for odd k , consider the graph G constructed by joining two complete graphs K_{2m-3} and K_{2m+3} with two independent edges e_1 and e_2 , where m is a sufficiently large integer. Then G is 2-connected, $G - E(C)$ is disconnected, $|G| = 4m$ and

$$\max\{d_G(x), d_G(y)\} \geq 2m + 2 = \frac{|G| + 4}{2}$$

for any two nonadjacent vertices x and y of $V(G)$. For any Hamiltonian cycle C , $G - E(C)$ has no $(k-2)$ -factor since C contains both e_1 and e_2 , and thus $G - E(C)$ consists of two components of odd order. Clearly, G satisfies all the conditions of the main theorem ([Theorem 6](#)) but has no k -factor containing a Hamiltonian cycle (by [Theorem 7](#)).

Acknowledgments

The authors would like to thank the referees for their extremely careful reading of the paper, and their detailed corrections and helpful suggestions. Without their help, this paper would never have had taken its present shape.

Y. Gao's work was supported by the Foundation for the distinguished young scholars of Shandong province (No. 2007BS01021). G. Li's work was supported by research grants from NSFC under grant numbers: 60673059, 10631070 and 60373025; and also by Taishan scholar fund from Shandong province.

References

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, North-Holland, Amsterdam, 1976.
- [2] G. Fan, New sufficient conditions for cycles in graphs, Journal of Combinatorial Theory, Series B 37 (1984) 221–227.
- [3] O. Ore, Note on Hamiltonian circuits, The American Mathematical Monthly 67 (1960) 55.
- [4] T. Iida, T. Nishimura, An Ore-Type condition for the existence of k -factors in graphs, Graphs and Combinatorics 7 (1991) 353–361.
- [5] T. Nishimura, A degree condition for the existence of k -factors, Journal of Graph Theory 16 (1992) 141–151.
- [6] H. Matsuda, Regular factors containing a given Hamiltonian cycle, Lecture Notes in Computer Science, Combinatorial Geometry and Graph Theory (2005) 123–130.
- [7] W.T. Tutte, The factors of graphs, Canadian Journal of Mathematics 4 (1952) 314–328.
- [8] W.T. Tutte, The factorization of linear graphs, Journal of London Mathematical Society 4 (1947) 107–111.